

Measurement of Dipolar Couplings in a Uniformly ^{13}C , ^{15}N -Labeled Membrane Protein: Distances between the Schiff Base and Aspartic Acids in the Active Site of Bacteriorhodopsin

Christopher P. Jaroniec,[†] Jonathan C. Lansing,[†]
Brett A. Tounge,^{†,§} Marina Belenky,[§] Judith Herzfeld,[§] and
Robert G. Griffin^{*†}

Department of Chemistry and
Francis Bitter Magnet Laboratory
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Department of Chemistry, Brandeis University
Waltham, Massachusetts 02254

Received August 23, 2001

In recent years, a number of magic-angle spinning (MAS) solid-state nuclear magnetic resonance (SSNMR) methods have been developed¹ for ^{13}C and ^{15}N resonance assignments in uniformly ^{13}C , ^{15}N -labeled peptides and proteins.² The ^{13}CO , $^{13}\text{C}\alpha$, $^{13}\text{C}\beta$, and amide ^{15}N chemical shifts can be used to estimate the backbone torsion angles ϕ and ψ .³ Additional constraints on the local structure in U- ^{13}C , ^{15}N -labeled systems can be obtained from measurements of the relative orientations of dipolar tensors.⁴ Long-range dipolar couplings in peptides and proteins can provide valuable information about tertiary structure. However, the accurate determination of weak dipolar interactions in U- ^{13}C , ^{15}N -labeled molecules is complicated by the presence of strong couplings.^{5,6} This problem can be circumvented by controlled “dilution” of the multiple spin system⁷ or by the use of spectrally selective dipolar recoupling techniques.^{8,9} To date, selective recoupling techniques have been applied to accurate measurements of multiple long-range ^{13}C – ^{13}C ¹⁰ and ^{13}C – ^{15}N ⁹ distances in small U- ^{13}C , ^{15}N -labeled peptides. In this Communication we demonstrate the application of frequency-selective REDOR (FSR)⁹ to the measurement of two ^{13}C – ^{15}N dipolar couplings in the active site of light-adapted [U- ^{13}C , ^{15}N]bacteriorhodopsin (bR) in its native purple membrane. The measured distances are in reasonable agreement with ones reported in recent diffraction structures of light-adapted bR, and the NMR methods described are directly applicable to bR photocycle intermediates, for which high-resolution diffraction structures are more difficult to obtain. The experiments presented here are the first example of long-range MAS NMR distance measurements in a U- ^{13}C , ^{15}N -labeled macromolecule.

Bacteriorhodopsin (bR) is a 26 kDa integral membrane protein produced by *Halobacterium salinarum*. The single polypeptide




Figure 1. Two-dimensional RFDR ^{13}C – ^{13}C chemical shift correlation spectrum of dark-adapted [U- ^{13}C , ^{15}N]bR displaying Asp C β –C γ and Glu C γ –C δ cross-peaks. Spectra were recorded at 11.7 T, 12.5 kHz MAS, and -80°C . To eliminate Asn C γ and Gln C δ signals in t_2 , the 2D RFDR sequence¹¹ was followed by a REDOR¹² ^{13}C – ^{15}N dipolar filter: CP– t_1 – $(\pi/2)_\psi$ –RFDR– $(\pi/2)$ –REDOR filter– t_2 . RFDR and REDOR filter lengths were 0.96 and 1.44 ms, respectively. Hypercomplex data were acquired by shifting phase ψ according to Ruben and co-workers.¹³ The data were acquired as (16, 512) complex points with dwell times (320, 20) μs . Each FID was 512 scans, with a 4.0-s recycle delay, resulting in a total measurement time of ~ 18 h. ^{13}C chemical shifts are indirectly referenced to the methyl ^1H resonance of DSS.¹⁴

chain forms a bundle of seven transmembrane helices enveloping a chromophore formed by a Schiff base (SB) between retinal and Lys216. Dark-adapted bR comprises two species: bR₅₅₅ and bR₅₆₈, with different retinal conformations. Light adaptation of bR, by irradiation with white light, converts bR₅₅₅ to bR₅₆₈, which is the starting point of the proton pumping photocycle (see ref 15 for a review on bR).

In the one-dimensional MAS spectrum of bR₅₆₈, the Schiff base nitrogen resonates ~ 135 ppm downfield from the ζ -NH₃⁺ groups of Lys residues and ~ 50 ppm downfield from the amide backbone peak.¹⁶ This enables the selective inversion of the SB nitrogen and FSR distance measurements⁹ to ^{13}C nuclei in the active site. Recent diffraction structures of bR₅₆₈ (from 1.55 to 2.9 Å resolution)^{17–19} report distances to the SB nitrogen in the 4.3–5.0 Å range for Asp85 C γ and in the 4.0–4.4 Å range for Asp212 C γ (see Supporting Information).

Figure 1 shows a region of a 2D RFDR¹¹ ^{13}C – ^{13}C chemical shift correlation spectrum of dark-adapted [U- ^{13}C , ^{15}N]bR displaying Asp and Glu side chain methylene to carboxyl cross-peaks.

(11) Bennett, A. E.; Ok, J. H.; Griffin, R. G.; Vega, S. *J. Chem. Phys.* **1992**, *96*, 8624.

(12) Gullion, T.; Schaefer, J. *J. Magn. Reson.* **1989**, *81*, 196.

(13) States, D. J.; Haberkorn, R. A.; Ruben, D. *J. Magn. Reson.* **1982**, *48*, 286.

(14) Wishart, D. S.; Bigam, C. G.; Yao, J.; Abildgaard, F.; Dyson, H. J.; Oldfield, E.; Markley, J. L.; Sykes, B. D. *J. Biomol. NMR* **1995**, *6*, 135.

(15) Haupts, U.; Tittor, J.; Oesterhelt, D. *Annu. Rev. Biophys. Biomol. Struct.* **1999**, *28*, 367.

(16) de Groot, H. J. M.; Harbison, G. S.; Herzfeld, J.; Griffin, R. G. *Biochemistry* **1989**, *28*, 3346.

(17) Essen, L. O.; Siebert, R.; Lehmann, W. D.; Oesterhelt, D. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95*, 11673.

(18) Belrhali, H.; Nollert, P.; Royant, A.; Menzel, C.; Rosenbusch, J. P.; Landau, E. M.; Pebay-Peyroula, E. *Structure* **1999**, *7*, 909.

(19) Luecke, H.; Schobert, B.; Richter, H. T.; Cartailler, J.; Lanyi, J. K. *J. Mol. Biol.* **1999**, *291*, 899.

* Correspondence author. E-mail: rgg@mit.edu.

[†] Massachusetts Institute of Technology.

[§] Brandeis University.

(1) Griffin, R. G. *Nature Struct. Biol.* **1998**, *5*, 508. Dusold, S.; Sebald, A. *Annu. Rep. Nucl. Magn. Reson. Spectrosc.* **2000**, *41*, 185.

(2) Rienstra, C. M.; Hohwy, M.; Hong, M.; Griffin, R. G. *J. Am. Chem. Soc.* **2000**, *122*, 10979. Balbach, J. J.; et al. *Biochemistry* **2000**, *39*, 13748. Detken, A.; et al. *J. Biomol. NMR* **2001**, *20*, 203. McDermott, A.; et al. *J. Biomol. NMR* **2000**, *16*, 209. Pauli, J.; Baldus, M.; van Rossum, B.; de Groot, H.; Oschkinat, H. *ChemBiochem* **2001**, *2*, 272.

(3) Spera, S.; Bax, A. *J. Am. Chem. Soc.* **1991**, *113*, 5490. Cornilescu, G.; Delaglio, F.; Bax, A. *J. Biomol. NMR* **1999**, *13*, 289.

(4) Feng, X.; Lee, Y. K.; Sandström, D.; Eden, M.; Maisel, H.; Sebald, A.; Levitt, M. H. *Chem. Phys. Lett.* **1996**, *257*, 314.

(5) Hodgkinson, P.; Emsley, L. *J. Magn. Reson.* **1999**, *139*, 46.

(6) Fyfe, C. A.; Lewis, A. R. *J. Phys. Chem. B* **2000**, *104*, 48.

(7) Reif, B.; Jaroniec, C. P.; Rienstra, C. M.; Hohwy, M.; Griffin, R. G. *J. Magn. Reson.* **2001**, *151*, 320.

(8) Raleigh, D. P.; Levitt, M. H.; Griffin, R. G. *Chem. Phys. Lett.* **1988**, *146*, 71. Takegoshi, K.; Nomura, K.; Terao, T. *Chem. Phys. Lett.* **1995**, *232*, 424.

(9) Jaroniec, C. P.; Tounge, B. A.; Herzfeld, J.; Griffin, R. G. *J. Am. Chem. Soc.* **2001**, *123*, 3507.

(10) Nomura, K.; Takegoshi, K.; Terao, T.; Uchida, K.; Kainosho, M. *J. Biomol. NMR* **2000**, *17*, 111.




Figure 2. Two-dimensional experiment for the measurement of dipolar couplings between the retinal SB nitrogen and Asp C γ in [U- ^{13}C , ^{15}N]-bR. Spectra were recorded at 7.4 T, 6494 Hz MAS, and -30°C . Light-adapted bR was prepared as described in the Supporting Information. Narrow and wide rectangles represent $\pi/2$ and π pulses, respectively. The ^{13}C Gaussian $3\pi/2$ pulse applied in the Asp C β spectral region following CP was 1.232 ms long. RFDR and REDOR filter times were both 1.232 ms. REDOR and FSR ^{15}N π pulse lengths were 13 μs (XY-4 phase cycling). During FSR (t_1) Gaussian π pulse lengths were 0.308 and 4.004 ms for Asp ^{13}C O and Lys216 $^{15}\text{N}\zeta$, respectively. Gaussian pulses were divided into 64 steps and truncated at 1%. The z -filters were $\Delta = 50 \mu\text{s}$. Proton decoupling was 125 and 100 kHz during mixing and acquisition, respectively (see ref 9). Phase cycle: $\phi_1 = 1$, $\phi_2 = 1$, $\phi_3 = 24$, $\phi_4 = 1$, $\phi_5 = 2$, $\phi_6 = 11111111$ 11111111 11111111 11111111 33333333 33333333 33333333 33333333, $\phi_7 = 11111111$ 22222222 33333333 44444444, $\phi_8 = 11223344$, $\phi_{rec'r} = 42244224$ 31133113 24422442 13311331 24422442 13311331 42244224 31133113, where 1 = x , 2 = y , 3 = $-x$, 4 = $-y$.

The resonance assignments are based on previous studies of selectively [$4\text{-}^{13}\text{C}$]Asp-labeled bR.²⁰ Asp85 and Asp212 C β -C γ cross-peaks are well-resolved, and, in principle, the distances to the SB can be measured via a 3D FSR experiment (where the decay rate of cross-peaks in the FSR dimension is determined by the dipolar coupling to SB ^{15}N). However, with the current technology, a 3D experiment for a membrane protein, such as bR, is subject to signal-to-noise and time limitations. Therefore, we have reduced the experiment to 2D (see Figure 2) by implementing a filter prior to FSR, designed to retain only Asp C γ resonances in the CO spectral region, while eliminating all other signals (although in practice some residual Glu C δ intensity remains). The filter consists of a selective storage pulse applied to Asp C β , followed by coherence transfer to C γ and C α via RFDR.¹¹ The subsequent REDOR filter eliminates signals in the CO spectral region, which originate from nuclei directly bonded to a ^{15}N nucleus (Asn C γ , Gln C δ , and C'). We have confirmed that the resulting 1D spectrum is equivalent to a weighted projection through the Asp C β -C γ and Glu C γ -C δ region in the 2D correlation spectrum (see Supporting Information).

FSR distance measurements from Lys216 N ζ to Asp85 C γ and Asp212 C γ in light-adapted bR are presented in Figure 3. Spectra in Figure 3a,b clearly show that Asp85 and Asp212 C γ are selectively dephased by the retinal SB ^{15}N , while the other Asp signals remain unchanged. FSR dephasing curves for several ^{13}C - ^{15}N distances in the 4.0–5.5 Å range were calculated using the SIMPSON NMR simulation software²¹ (see Supporting Information). Experimental S/S_0 curves for Asp85 and Asp212 are




Figure 3. Distance measurements from the Schiff base (Lys216 N ζ) to Asp85 C γ and Asp212 C γ in light-adapted [U- ^{13}C , ^{15}N]bR. (a) Reference (S_0) and (b) REDOR (S) spectra recorded using the pulse sequence in Figure 2 with $t_1 = 19.712$ ms. Experimental spectra (O) and fits with Gaussian line shapes (lines) are juxtaposed. The deviation of the baseline from zero in (a) and (b) is within the experimental noise level as determined from the entire spectrum. (c) Asp85 C γ and (d) Asp212 C γ REDOR S/S_0 curves. Experimental data (O) were obtained with 32,768 and 65,536 transients for $t_1 < 15$ ms and $t_1 > 15$ ms, respectively (interleaving S_0 and S spectra for each t_1 point in blocks of 256 transients). The error bars in the experimental S/S_0 points were calculated from the uncertainties in the Gaussian line shape fits. Using a 2.0-s recycle delay, the total duration of the experiment was ~ 10 days. Simulations (lines) for 4.0, 4.5, 5.0, and 5.5 Å carbon–nitrogen distances were generated using SIMPSON.²¹ The best-fit distances are 4.7 ± 0.3 Å for Asp85 and 4.9 ± 0.5 Å for Asp212.

compared with 4.0, 4.5, 5.0, and 5.5 Å simulations in Figure 3c,d. Asp85 and Asp212 distances were found to be 4.7 ± 0.3 and 4.9 ± 0.5 Å, respectively. The Asp85 distance is in agreement with recent diffraction distances (4.3–5.0 Å), and the Asp212 distance appears to be somewhat longer than the diffraction values (4.0–4.4 Å).^{17–19}

In summary, we have applied frequency-selective REDOR to distance measurements between the Schiff base and aspartic acids in the active site of uniformly ^{13}C , ^{15}N -labeled bacteriorhodopsin. The measured distances are in reasonable agreement with recent diffraction studies and complement previous structural investigations of the bR active site by SSNMR.²² The methods described can be applied to the characterization of bR photocycle intermediates.

Acknowledgment. This research was supported by NIH Grants GM-23289, GM-36810, and RR-00995. C.P.J. thanks the NSF for a Predoctoral Fellowship, and B.A.T. thanks the American Cancer Society for a Postdoctoral Fellowship (PF-99-260-01-GMC).

Supporting Information Available: Table of diffraction distances, bR spectra, [U- ^{13}C , ^{15}N]bR preparation, and simulation details (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA016923R

(21) Bak, M.; Rasmussen, J. T.; Nielsen, N. C. *J. Magn. Reson.* **2000**, *147*, 296.

(22) Griffiths, J. M.; Bennett, A. E.; Engelhard, M.; Siebert, F.; Raap, J.; Lugtenburg, J.; Herzfeld, J.; Griffin, R. G. *Biochemistry* **2000**, *39*, 362.

(20) Metz, G.; Siebert, F.; Engelhard, M. *Biochemistry* **1992**, *31*, 455. Metz, G.; Siebert, F.; Engelhard, M. *FEBS Lett.* **1992**, *303*, 237.

Supporting Information

Christopher P. Jaroniec, Jonathan C. Lansing, Brett A. Tounge, Marina Belenky,
Judith Herzfeld, and Robert G. Griffin

**Measurement of Dipolar Couplings in a Uniformly ^{13}C , ^{15}N Labeled
Membrane Protein: Distances Between the Schiff Base and Aspartic
Acids in the Active Site of Bacteriorhodopsin**

Journal of the American Chemical Society

Preparation of light-adapted [U-¹³C,¹⁵N]bacteriorhodopsin

Uniformly ¹³C,¹⁵N-labeled peptone for the culture medium of *Halobacterium salinarum*¹ was obtained from the anaerobic acid hydrolysis of *Methylophilus methylotrophus* grown on ¹³C-labeled methanol and ¹⁵N-labeled ammonium sulfate.² The purple membranes were isolated using the method of Oesterhelt and Stoeckenius.³ Prior to the NMR experiments the purple membrane sample (~40 mg) was washed 3 times with a 0.1 M NaCl solution at pH 10.0. Following each wash the sample was centrifuged for 1 h at ~43000g. After the final spin the bR pellet was packed into a transparent single-crystal sapphire rotor (5 mm o.d.) and sealed.

The rotor was placed in the NMR probe and the light-adapted state (bR₅₆₈) was prepared *in situ* via overnight irradiation with white-light at 0 °C (the light from a 1000 W xenon lamp was filtered through water and heat absorbing filter and delivered to the sample via a glass optic fiber bundle). Subsequently, the sample temperature was decreased to ca. -30 °C. The presence of pure bR₅₆₈ was confirmed prior to and following the distance measurements by monitoring the characteristic chemical shift of the retinal Schiff base nitrogen in a 1D ¹⁵N CPMAS spectrum.⁴

¹ Gochauer, M. B.; Kushner, D. *Can. J. Microbiol.* **1969**, *15*, 1157-1165.

² Batey, R. T.; Kujawinski, E.; Puglisi, J. D.; Williamson, J. R. *Nucleic Acid Res.* **1992**, *20*, 4515-4523. Batey, R. T.; Williamson, J. R. *Methods Enzymol.* **1995**, *261*, 4515-4523.

³ Oesterhelt, D.; Stoeckenius, W. *Methods Enzymol.* **1974**, *31*, 667-678.

⁴ de Groot, H. J. M.; Harbison, G. S.; Herzfeld, J.; Griffin, R. G. *Biochemistry* **1989**, *28*, 3346-3353.

Numerical Simulations

Numerical simulations of the FSR experiment¹ in a heteronuclear two-spin system were performed using the SIMPSON NMR simulation software.² The simulations included the exact profiles of the Gaussian pulses. In the FSR experiment on bR, the dephasing of the Asp85 and Asp212 C γ signals is primarily a function of the dipolar coupling to the retinal Schiff base nitrogen; the couplings from C γ to other ¹⁵N, as well as Asp C γ -C β , etc. J-couplings are suppressed.¹ The dephasing is also affected by the non-negligible magnitudes of the ¹³C and ¹⁵N anisotropic chemical shielding (CSA) interactions, and to a minor extent their relative orientations.³ Therefore, ¹⁵N and ¹³C CSAs were included in the simulations (previously reported bR₅₆₈ ¹⁵N and ¹³C CSA principal values were used^{4,5}). Relative orientations of the dipolar and CSA tensors are not known for bR, and simulations were performed for 1024 sets of different relative tensor orientations. Subsequently, the average dephasing curves expected for different distances in the 4.0-5.5 Å regime were calculated, and used to extract the internuclear distances using standard reduced χ^2 analysis.⁶ We found that for each value of the dipolar coupling the simulations for different relative tensor orientations are clustered very closely around the average dephasing curve, with the largest deviation observed when the principal axis systems of the three tensors are aligned. The systematic errors in the estimated dipolar coupling introduced by the choice of an incorrect relative tensor orientation could be ignored, since they were significantly smaller (< 0.1 Å) than the random errors in the measured distances (ca. 0.3-0.5 Å).

¹ Jaroniec, C. P.; Tounge, B. A.; Herzfeld, J.; Griffin, R. G. *J. Am. Chem. Soc.* **2001**, *123*, 3507-3519.


² Bak, M.; Rasmussen, J. T.; Nielsen, N. C. *J. Magn. Reson.* **2000**, *147*, 296-330.

³ Additional simulations indicate that CSA dependence is significantly reduced when higher MAS frequencies ($\omega_r \gg |\omega_{CSA}|$) are employed. In this work the maximum spinning frequencies were limited to ~6.5 kHz, since we have used the low-temperature NMR probe equipped with a 5-mm stator, capable of trapping and characterizing bR photocycle intermediates *in situ*.


⁴ de Groot, H. J. M.; Harbison, G. S.; Herzfeld, J.; Griffin, R. G. *Biochemistry* **1989**, *28*, 3346-3353.

⁵ Metz, G.; Siebert, F.; Engelhard, M. *Biochemistry* **1992**, *31*, 455-462.

⁶ Shoemaker, D. P.; Garland, C. W.; Nibler, J. W. *Experiments in Physical Chemistry*; McGraw-Hill: New York, 1989 and references therein.



Supporting Information Figure 1. 2D RFDR ^{13}C - ^{13}C chemical shift correlation spectrum of dark-adapted [$\text{U-}^{13}\text{C}, ^{15}\text{N}$]bacteriorhodopsin displaying Asp C β -C γ and Glu C γ -C δ cross-peaks (a) and filtered 1D ^{13}C spectrum of bR containing only Asp C γ and Glu C δ resonances in the CO spectral region (b). The FID in (b) consisted of 16384 scans, with a 4.0-s recycle delay, yielding a total measurement time of ~ 18 h. Other experimental details are described in Figure 1 caption and text.



Supporting Information Figure 2. Illustration of the active site of bR. Shown are the B, C, F and G α -helices, the chromophore (comprising retinal with its Schiff base linkage to Lys216), and the Asp 85 and Asp 212 sidechains. The distances between the Schiff base nitrogen and Asp C γ , measured in this work, are indicated by dotted lines.

Supporting Information Table 1. Diffraction Distances from the Retinal Schiff Base Nitrogen in bR₅₆₈.

PDB	Resolution (Å)	to Asp85 C γ (Å)	to Asp212 C γ (Å)
1BRR ¹	2.9	5.06	4.14
		5.01	4.13
		4.96	3.98
1QHJ ²	1.9	4.28	4.41
1C3W ³	1.55	4.45	4.12

¹ Essen, L. O.; Siebert, R.; Lehmann, W. D.; Oesterhelt, D. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 11673-11678.

² Belrhali, H.; Nollert, P.; Royant, A.; Menzel, C.; Rosenbusch, J. P.; Landau, E. M.; Pebay-Peyroula, E. *Structure* **1999**, *7*, 909-917.

³ Luecke, H.; Schobert, B.; Richter, H. T.; Cartailler, J.; Lanyi, J. K. *J. Mol. Biol.* **1999**, *291*, 899-911.